

REMARKS ON GRAVITY AND QUANTUM GEOMETRY

ROBERT CARROLL
UNIVERSITY OF ILLINOIS, URBANA, IL 61801

ABSTRACT. Some relations between conformal relativity and Weyl-Dirac theory are rephrased and clarified.

1. BACKGROUND

In recent times, stimulated by seminal paper of E. Verlinde [60], there has been a deluge of material about entropy and gravity (we refer to Padmanabhan [42, 43, 44] for background and some references to the new material along with some striking new results on equipartition). The strong connections of thermodynamics, entropy, and gravity have been in the air for some time, going back to Beckenstein, Jacobson, and Padmanabhan (cf. [6, 30, 42]) and to others too numerous to name. It is also well known that quantum mechanics is related to entropy (and information theory) and we cite here only a recent paper by Caticha [22] which summarizes much research in this connection (going back to Nagasawa, Nelson, and Smolin among many others - see e.g. [36, 37, 57]). In [12, 13, 14, 15, 16, 17, 18] we have also described connections between quantum mechanics, Weyl geometry, and Weyl-Dirac quantum theory following [1, 3, 4, 5, 8, 20, 21, 24, 27, 28, 29, 38, 39, 51, 53, 54, 55, 63] (note that publication information can often be found on the electronic bulletin board entries which we often use in our referencing). The key idea here, going back at least to Audretsch et al and Santamato, is that quantum mechanics is naturally associated with a Weyl geometry, and hence to gravitational ideas. Another manifestation of gravity-quantum connections appears in papers of Mannheim and Bender (see e.g. [7, 34]) where one refers e.g. to intrinsically quantum mechanical gravity. This background of quantum mechanics in geometry (or gravity) says nothing about “quantum gravity”, as one imagines might be built up in various manners (see e.g. [31, 32, 35, 41, 50, 58, 61, 62]). Rather the general motivating idea seems to be in the spirit of accepting a universe guided by information theory plus some ingredients (space, time, matter) and describing how everything fits together. Never mind trying

Date: July, 2010.
email: rcarroll@math.uiuc.edu.

to build a universe from scratch. Of course this encompasses also trying to understand the microstructure of spacetime, the cosmological constant, and other matters.

2. WEYL DIRAC THEORY

We review first two derivations of Dirac-Weyl theory following [27]. The first follows Dirac in avoiding Proca terms and the second involves integrable Weyl geometry. Thus let $ds^2 = g_{ab}dx^a dx^b$ be based on a Riemannian metric g_{ab} with Levi-Civita connection coefficients (Christoffel symbols)

$$(2.1) \quad \Gamma_{\mu\nu}^\lambda = \frac{1}{2}g^{\lambda\sigma}(\partial_\nu g_{\mu\sigma} + \partial_\mu g_{\nu\sigma} - \partial_\sigma g_{\mu\nu})$$

Recall also that a comma $(,n)$ denotes ∂_n and a semicolon $(;n)$ denotes a covariant derivative ∇_n so that

$$(2.2) \quad B_{;\nu}^\mu = \nabla_\nu B^\mu = \partial_\nu B^\mu + B^\sigma \Gamma_{\sigma\nu}^\mu; \quad B_{\mu;\nu} = \nabla_\nu B_\mu = \partial_\nu B_\mu - B_\sigma \Gamma_{\mu\nu}^\sigma$$

If a vector is parallel transported x^ν to $x^\nu + dx^\nu$ one has via (2.2) **(2A)** $dB = 0$ where $B = (g_{\mu\nu}B^\mu B^\nu)^{1/2}$ and one checks that **(2B)** $\nabla g_{\mu\nu} = 0 = \nabla g^{\mu\nu}$. The (Riemann-Christoffel) curvature tensor is

$$(2.3) \quad R_{\sigma\mu\nu}^\lambda = -\partial_\nu \Gamma_{\sigma\mu}^\lambda + \partial_\mu \Gamma_{\sigma\nu}^\lambda - \Gamma_{\sigma\mu}^\alpha \Gamma_{\alpha\nu}^\lambda + \Gamma_{\sigma\nu}^\alpha \Gamma_{\alpha\mu}^\lambda$$

and if e.g. B^μ is parallel transported around an infinitesimal closed parallelogram the total change is **(2C)** $\Delta B^\lambda = B^\sigma R_{\sigma\mu\nu}^\lambda dx^\mu \delta x^\nu$ (whereas $\Delta B = 0$).

Now for Weyl geometry both the length and direction of a vector change under parallel transport and one has a (symmetric) Weyl connection $\hat{\Gamma}_{\mu\nu}^\lambda = \hat{\Gamma}_{\nu\mu}^\lambda$ with e.g.

$$(2.4) \quad dB^\mu = -B^\sigma \hat{\Gamma}_{\sigma\nu}^\mu dx^\nu; \quad dB = B w_\nu dx^\nu$$

where w_ν is called a Weyl vector. In order to be compatible with the a given metric g_{ab} one must have now

$$(2.5) \quad \hat{\Gamma}_{\mu\nu}^\lambda = \Gamma_{\mu\nu}^\lambda + g_{\mu\nu}w^\lambda - \delta_\nu^\lambda w_\mu - \delta_\mu^\lambda w_\nu$$

In particular (denoting Weyl covariant derivatives via a colon $(:)$ (as well as a hat) it follows that **(2D)** $g_{:\lambda}^{\mu\nu} = \hat{\nabla}_\lambda g^{\mu\nu} = -2g^{\mu\nu}w_\lambda$. One assumes $\hat{\Gamma}_{\mu\nu}^\lambda = \hat{\Gamma}_{\nu\mu}^\lambda$ and formulas analogous to (2.2) hold for $\hat{\Gamma}_{\mu\nu}^\lambda$. Now however parallel transport around an infinitesimal closed parallelogram yields **(2E)** $\Delta B = BW_{\mu\nu}dx^\mu \delta x^\nu$ where **(2F)** $W_{\mu\nu} = \partial_\nu w_\mu - \partial_\mu w_\nu$. In case $w_\lambda = \partial_\lambda \phi$ one has an integrable Weyl geometry with $W_{\mu\nu} = 0$ and $\Delta B = 0$.

Let now a vector of length B transported around a closed loop to achieve a new length **(2G)** $B_{new} = B + \int_S BW_{\mu\nu}dS^{\mu\nu}$ with S the area enclosed by the loop and $dS^{\mu\nu}$ an area element. In this connection one considers local

Weyl gauge transformations (WGT) **(2H)** $B \rightarrow \tilde{B} = \exp(\lambda)B$ where $\lambda(x^\nu)$ is a differentiable function leading to

$$(2.6) \quad g_{\mu\nu} \rightarrow \tilde{g}_{\mu\nu} = e^{2\lambda} g_{\mu\nu}; \quad g^{\mu\nu} \rightarrow \tilde{g}^{\mu\nu} = e^{-2\lambda} g^{\mu\nu}$$

This is all compatible with Weyl gauge transformations, namely **(2I)** $w_\nu \rightarrow \tilde{w}_\nu = w_\nu + \partial_\nu \lambda$ and a Weyl weight (or power) $n = \Pi(\psi)$ is defined for quantities transforming via **(2J)** $\psi \rightarrow \tilde{\psi} = \exp(n\lambda)\psi$ (cf. [10, 27, 48] for more discussion). In particular **(2K)** $\hat{\Gamma}_{\mu\nu}^\lambda \rightarrow \hat{\Gamma}_{\mu\nu}^\lambda$ with $\Pi(\hat{\Gamma}_{\mu\nu}^\lambda) = 0$, $\Pi(g_{\mu\nu}) = 2$, and $\Pi(\sqrt{-g}) = 4$, etc.

Now in the Dirac formulation of a combined relativity plus electromagnetism (EM) one arrives at an action (cf. [27])

$$(2.7) \quad I = \int \left[W^{\lambda\mu} W_{\lambda\mu} - \beta^2 R + \sigma \beta^2 |\mathbf{w}|^2 + (\sigma + 6)(\nabla\beta)^2 + \right. \\ \left. + 2\sigma\beta w^\lambda \partial_\lambda \beta + 2\Lambda\beta^4 + L_M \right] \sqrt{-g} d^4x$$

Then in order to avoid Proca terms Dirac set $\sigma = 0$ to arrive at

$$(2.8) \quad I = \int \left[W^{\lambda\mu} W_{\lambda\mu} - \beta^2 R + 6(\nabla\beta)^2 + 2\Lambda\beta^4 + L_M \right] \sqrt{-g} d^4x$$

There is however another approach due to Rosen [48] via an integrable Weyl-Dirac theory in which the action is given by (2.9) below and σ can take arbitrary values (cf. [27] from which we extract). Thus take $w_\nu = \partial_\nu w$ (so $W_{\mu\nu} = 0$) and set **(2L)** $b_\mu = c\partial_\mu[\log(\beta)]$ with $W = w + b$. In fact we could now take $w = 0$ and still have an integrable Weyl geometry with $W_\nu = c\partial_\nu \log(\beta)$ (note this is not necessary however). The action becomes then

$$(2.9) \quad I = \int \sqrt{-g} d^4x \left[-\beta^2 R + (\sigma + 6)(\nabla\beta)^2 + 2\Lambda\beta^4 + L_M \right]$$

(see Remark 2.1 below).

In [14, 15, 16, 18] we considered a conformal map from an Einstein frame action **(2M)** $S_{GR} = \int d^4x \sqrt{-g} [R - \alpha(\nabla\psi)^2 + 16\pi L_M]$ to a Brans-Dicke (BD) form called conformal GR, or CGR, which we specify now after deleting the L_M and Λ terms, so that CGR is represented via

$$(2.10) \quad \hat{S}_{GR} = \int d^4x \sqrt{-g} e^{-\psi} \left[\hat{R} - \left(\alpha - \frac{3}{2} \right) (\hat{\nabla}\psi)^2 \right] = \\ = \int d^4x \sqrt{-\hat{g}} \left[\hat{\phi} \hat{R} - \left(\alpha - \frac{3}{2} \right) \frac{(\hat{\nabla}\hat{\phi})^2}{\hat{\phi}} \right]$$

(here $\Omega^2 = \phi = \exp(\psi) = \hat{\phi}^{-1}$). This is an integrable Weyl geometry based originally on $\hat{g}_{ab} = \Omega^2 g_{ab}$, $\Omega^2 = \exp(\psi) = \phi = \hat{\phi}^{-1}$, $w_\alpha = \partial_\alpha \psi =$

$\partial_\alpha \hat{\phi}/\hat{\phi}$ and a conformal mass $\hat{m} = \hat{\phi}^{-1/2}m$ (cf. [8]). Further $\hat{m}$ can be identified with a quantum mass field $\mathfrak{M} \sim \beta$ from a Bohmian Weyl-Dirac theory with a quantum potential Q determined via $\mathfrak{M}^2 = m^2 \exp(Q)$ (cf. [8, 12, 13, 14, 53]). Thus $\mathfrak{M} = \beta = \hat{m} = \hat{\phi}^{-1/2}m$ and $\Omega^2 = \hat{\phi}^{-1} = \mathfrak{M}^2/m^2 = \beta^2/m^2$. Now from (2L) for example, given w_α already of the form $w_\alpha = -\partial_\alpha \psi = \partial_\alpha \hat{\phi}/\hat{\phi}$, with $\beta = m\hat{\phi}^{-1/2}$, we find that $w_\alpha = -2\partial_\alpha \log(\beta)$. This means that one could take $w = 0$ in $W = w + b$ with $b_\alpha = c\partial_\alpha \log(\beta)$ in comparing Weyl-Dirac actions.

REMARK 2.1. Note that, given $w_\alpha = -2\partial_\alpha \log(\beta)$, if we choose $b_\alpha = c\partial_\alpha \log(\beta)$ with $c \neq 2$, then $W_\alpha = w_\alpha + b_\alpha = (c-2)\partial_\alpha \log(\beta)$ so $W_{\mu\nu} = 0$. However no such manipulation is needed since the unadulterated Weyl-Dirac integrand from [27] (p. 116) is $(-\beta^2 \sim \sum(\partial_j \beta)^2 \sim (\nabla \beta)^2 \sim (\partial \beta)^2$ and $|\mathbf{w}|^2 = 4(\partial \beta)^2/\beta^2$)

$$(2.11) \quad \left[-\beta^2 R + \sigma \beta^2 |\mathbf{w}|^2 + 2\sigma \beta w^\lambda \partial_\lambda \beta + (\sigma + 6)(\partial \beta)^2 + 2\Lambda \beta^4 + L_M \right]$$

Here $|\mathbf{w}|^2 = 4(\partial \beta/\beta)^2$ so $\beta^2 |\mathbf{w}|^2 = 4(\partial \beta)^2$ and $\beta w^\lambda \partial_\lambda \beta = -2(\partial \beta)^2$ so, omitting the Λ and L_M terms, the integrand is

$$(2.12) \quad -\beta^2 R + 4\sigma(\partial \beta)^2 - 4\sigma(\partial \beta)^2 + (\sigma + 6)(\partial \beta)^2 = -\beta^2 R + (\sigma + 6)(\partial \beta)^2$$

One sees that the $4\sigma(\partial \beta)^2$ terms cancel. ■

Now, following [14], for (2.10) we note first from [14] that (2N) $\hat{R}\hat{\phi}\sqrt{-\hat{g}} = \hat{R}\hat{\phi}\sqrt{-g} = (\beta^2/m^2)\hat{R}\sqrt{-g}$ and (2O) $\beta^2 \hat{R} = \beta^2 R - 6\beta^2 \nabla_\lambda w^\lambda + 6\beta^2 |\mathbf{w}|^2$. Hence the first integral in (2.10) takes the form

$$(2.13) \quad \begin{aligned} I_1 &= \int d^4x \sqrt{-\hat{g}} \hat{\phi} \hat{R} = \int d^4x \sqrt{-g} \phi \hat{R} = \int d^4x \sqrt{-g} \hat{R} \frac{\beta^2}{m^2} = \\ &= \frac{1}{m^2} \int d^4x \sqrt{-g} \left[\beta^2 R - 6\beta^2 \nabla_\lambda w^\lambda + 6\beta^2 |\mathbf{w}|^2 \right] \end{aligned}$$

and $-\beta^2 \nabla_\lambda w^\lambda = -\nabla_\lambda (\beta^2 w^\lambda) + 2\beta \partial_\lambda w^\lambda$ so the divergence term vanishes upon integration and (2.13) becomes

$$(2.14) \quad \begin{aligned} I_1 &= \frac{1}{m^2} \int d^4x \sqrt{-g} \left[\beta^2 R + 12\beta(\partial_\lambda \beta) w^\lambda + 6\beta^2 |\mathbf{w}|^2 \right] = \\ &= \frac{1}{m^2} \int d^4x \sqrt{-g} \left[\beta^2 R + 12(\beta \partial_\lambda \beta) \left(\frac{-2\partial^\lambda \beta}{\beta} \right) + 6\beta^2 \left(\frac{2\partial \beta}{\beta} \right)^2 + \right. \\ &\quad \left. + 6\beta^2 \left(\frac{2\partial \beta}{\beta} \right)^2 \right] = \frac{1}{m^2} \int d^4x \sqrt{-g} \beta^2 R \end{aligned}$$

The second integral in (2.10) is now $[\gamma = \alpha - (3/2)]$

$$(2.15) \quad \begin{aligned} I_2 &= -\gamma \int d^4x \sqrt{-\hat{g}} \frac{(\partial \hat{\phi})^2}{(\hat{\phi})^2} \hat{\phi} = \\ &= -\gamma \int d^4x \sqrt{-g} \phi \left(\frac{-2\partial\beta}{\beta} \right)^2 = -\frac{4\gamma}{m^2} \int d^4x \sqrt{-g} (\partial\beta)^2 \end{aligned}$$

Consequently (2.10) becomes

$$(2.16) \quad \hat{S}_{GR} = I_1 + I_2 = \frac{1}{m^2} \int d^4x \sqrt{-g} \left[\beta^2 R - 4 \left(\alpha - \frac{3}{2} \right) (\partial\beta)^2 \right]$$

This will then represent an integrable Weyl-Dirac theory provided that e.g. (2.16) $\equiv$ (2.12) which means that (modulo a factor of $(1/m^2)$) we choose $\sigma + 6 = -4[\alpha - (3/2)]$ or $\sigma = -4\alpha$.

REMARK 2.2 We note that there is a typo in equation (6.7), p. 59 (and in (3.33), p. 237) in [14] namely $3 - 4\alpha$ should be $6 - 4\alpha$. Also on pp. 58-59 and 236-237 one should have $\phi = \exp(\psi) = \hat{\phi}^{-1} = \Omega^2 = (\beta/m)^2 = (\mathfrak{M}/m)^2$ consistently and L_M should be deleted in (6.6)-(6.7), p. 59 and (3.32)-(3.33), p. 237. A multiplier $\hat{\phi}^2$ should also be added to **(6C)** on p. 59 and to **(★)** on p. 237. In [18], on p. 155, line -7, one should change ψ to $-\psi$, and note that the choice of comparison actions on p. 157, equation (52), was too constrained; we should have used the integrable Weyl-Dirac action (2.9) above with arbitrary σ . Also in [18], p. 157, equation (51) should have $6 - 4\alpha$ and on p. 167, line 13, insert: $\dots u$ in **(●●)** denotes $\dots$. ■

REMARK 2.3. We mention here a related discussion in [40] of quantum mechanics in connection with Weyl geometry and the Palatini approach to gravitation. This is connected to conformal aspects of the Palatini approach (as in [2, 11, 49]). We mention also [33] for a related treatment of geometry and the quantum. ■

REFERENCES

- [1] R. Adler, M. Bazin, and M. Schiffer, Introduction to general relativity, McGraw-Hill, 1975
- [2] G. Allemandi, M. Capone, S. Capozziello, and M. Francaviglia, hep-th 0409148
- [3] J. Audretsch, Phys. Rev. D, 24 (1981), 1470-1477; 27 (1983), 2872-2884
- [4] J. Audretsch, F. Gähler, and N. Straumann, Comm. Math. Phys., 95 (1984), 41-51
- [5] J. Audretsch and C. Lämmerzahl, Class. Quant. Gravity, 5 (1988), 1285-1295
- [6] J. Beckenstein, Phys. Rev. D, 7 (1973), 2333
- [7] C. Bender and P. Mannheim, Phys. Rev. D, 78 (2008), 025022; Phys. Rev. Lett., 100 (2008), 110402; hep-th 0807.2607
- [8] R. Bonal, I. Quiros, and R. Cardenas, gr-qc 0010010
- [9] C. Brans and R. Dicke, Phys. Rev., 124 (1961), 925-935
- [10] V. Canuto, P. Adams, S. Hsieh, and E. Tsiang, Phys. Rev. D, 16 (1977), 1643-1663
- [11] S. Capozziello, F. Darabi, and D. Vernieri, gr-qc 1006.0454
- [12] R. Carroll, Fluctuations, information, gravity, and the quantum potential, Springer, 2006
- [13] R. Carroll, On the quantum potential, Arima Publ., 2007
- [14] R. Carroll, On the emergence theme of physics, World Scientific, 2010
- [15] R. Carroll, Progress in Physics, 4 (2007), 22-24; 1 (2008), 21-24 and 2 (2008), 89-90
- [16] R. Carroll, math-ph 0701077, 0703065, 0705.3921, 0712.3251; 0507027
- [17] R. Carroll, Foundations of Physics, 35 (2005), 131-154
- [18] R. Carroll, Quantum Potential as Information: A mathematical survey, To appear in a book on quantum information science, Editors: Felloni, Singh, Licata, and Sakaji
- [19] R. Carroll, Theor. and Math. Phys., 152 (2007), 904-914
- [20] C. Castro, Found. Phys., 22 (1992), 569-615; 37 (2007), 366-409; Found. Phys. Lett., 4 (1991), 81-99; Jour. Math. Phys., 31 (1990), 2633-2626
- [21] C. Castro and J. Mahecha, Progress in Physics, 1 (2006), 38-45
- [22] A. Caticha, quant-ph 0907.4335 and 1005.2357
- [23] R. Dicke, Phys. Rev., 125 (1962), 2163-2167
- [24] P. Dirac, Proc. Royal Soc. London A, 209 (1951), 291-296; 212 (1952), 330-339; 332 (1973), 403-418, and 338 (1974), 439-446
- [25] V. Faraoni, Cosmology in scalar tensor gravity, Kluwer, 2004
- [26] O. Gron and S. Hervik, Einstein's general theory of relativity, Springer, 2007
- [27] M. Israelit, The Weyl-Dirac theory and our universe, Nova Science Pub., 1999; Found. Phys. 29 (1999), 1303-1322; 32 (2002), 295-321 and 945-961; 28 (1998), 205-228; 29 (1999), 1303-1322, 35 (2005), 1725-1748, and 1769-1782, 37 (2007), 1628-1642
- [28] M. Israelit, Gen. Relativ. Grav., 29 (1997), 1411-1429 and 1597-1614; gr-qc 0710.3410, 0710.3690, 0710.3913, 07103923, and 0905.2482
- [29] M. Israelit and N. Rosen, Found. Phys., 13 (1983), 1023-1045
- [30] T. Jacobson, Phys. Rev. Lett., 75 (1995), 1260
- [31] C. Kiefer, Quantum gravity, Oxford Univ. Press, 2004
- [32] F. Klinkhamer and G. Volovik, gr-qc 0403037, 0503090, 0711.3170, 0806.2805, 0807.3896, 0808.3896, and 0811.4347; astro-ph 0905.1419; hep-th 0907.4887
- [33] B. Koch, gr-qc 1004.2879 and 1004.3240
- [34] P. Mannheim, hep-th 0707.2283, 0809.1200, 0909.0212, 0912.2635, and 1005.5108; astro-ph 0505266

- [35] F. Markopoulou, gr-qc 0909.1861, 07070907, and 9811053
- [36] M Nagasawa, Schrödinger equation and diffusion theory, Birkhäuser, 1997; Stochastic processes in quantum physics, Birkhäuser, 2000
- [37] E. Nelson, Quantum fluctuations, Princeton Univ. Press, 1985
- [38] H. Nikolić, quant-ph 0512065, 0602024, 0603207, 0610138, 0703071, 0804.4564, 0811.1905; hep-th 9798.3962, 0605250, 0702069, 0705.3542, 0708.0729, 0806.1431, 0801.4471, 0904.2287; gr-qc 0111029, 0312063, 0805.2555, 0909035
- [39] J. Noldus, gr-qc 0508104
- [40] M. Novello, J. Salim, and F. Falciano, gr-qc 0901.3741
- [41] D. Oriti, Approaches to quantum gravity, Cambridge Univ. Press, 2009
- [42] T. Padmanabhan, Gravitation, Cambridge Univ. Press, 2010
- [43] T. Padmanabhan, gr-qc 0912.3165, 0911.5004, and 1003.5665
- [44] T. Padmanabhan, Class. Quant. Grav., 21 (2004), 4485; gr-qc 0705.2533, 0903.154, 0910.0839, and 0911.1403; astro-ph 0603114
- [45] T. Padmanabhan and A. Paranjape, gr-qc 0701003
- [46] I. Quiros, gr-qc 9904004 and 9905071; hep-th 0009169
- [47] I. Quiros, R. Bonal, and R. Cardenas, gr-qc 998075 and 0002071
- [48] N. Rosen, Found. Phys., 12 (1982), 213-248
- [49] E. Rosenthal, gr-qc 0809.2953
- [50] C. Rovelli, Quantum gravity, Cambridge Univ. Press, 2006
- [51] E. Santamato, Phys. Rev., 29 (1984), 216-222; 32 (1985), 2615-2621; Jour. Math. Phys., 25 (1984), 2477-2480; Phys. Lett. A, 130 (1988), 199-202
- [52] E. Scholz, astro-ph 0409635; gr-qc 0511113
- [53] F. and A. Shojai, gr-qc 0306099, 0404102 and 0409053; Class. Quant. Grav., 21 (2004), 1-9
- [54] F. and A. Shojai and M. Golshani, gr-qc 9903049 and 9903050
- [55] F. Shojai and M. Golshani, gr-qc 9903047
- [56] F. Shojai and S. Molladavoudi, gr-qc 0708.0620
- [57] L. Smolin, Phys. Lett. A, 113 (1986), 347; quant-ph 0609109
- [58] T. Thiemann, Modern canonical quantum general relativity, Cambridge Univ. Press, 2007
- [59] S. Tiwari, Phys. Lett. A, 135 (1989), 315-319; physics 0405136, 0408053, and 0502035; quant-ph 0109048, 0606031, and 0612139; gr-qc 0307079
- [60] E. Verlinde, hep-th 1001.0785
- [61] G. Volovik, The universe in a helium droplet, Oxford Univ. Press, 2003
- [62] X. Wen, QFT of many body systems, Oxford Univ. Press, 2004 2004
- [63] J. Wheeler, Phys. Rev. D, 41 (1990), 431-441; hep-th 9706215, 9708088, 0002068, and 0305017; gr-qc 9411030